

## PAPER NAME

**Manuscript Syarat Khusus.pdf**

---

## WORD COUNT

**6603 Words**

## CHARACTER COUNT

**36009 Characters**

## PAGE COUNT

**20 Pages**

## FILE SIZE

**479.4KB**

## SUBMISSION DATE

**Dec 15, 2023 2:36 PM GMT+7**

## REPORT DATE

**Dec 15, 2023 2:37 PM GMT+7**

---

**● 6% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 6% Publications database
- Crossref Posted Content database
- Crossref database

**● Excluded from Similarity Report**

- Internet database
- Quoted material
- Small Matches (Less than 8 words)
- Bibliographic material
- Cited material

# RELATIONSHIP BETWEEN ABUNDANCE OF *Chlamydomonas* spp AND *Chlorella* spp ON CLINICAL PERFORMANCE OF RED TILAPIA (*Oreochromis niloticus*) IN SILVOFISHERY PONDS

Linayati Linayati<sup>1</sup>, Nguyen Huu Yen Nhi<sup>2</sup>, Heri Ariadi<sup>1\*</sup>, Tri Yusufi Mardiana<sup>1</sup>, Ashari Fahrurrozi<sup>1</sup>, M. Bahrus Syakirin<sup>1</sup>

<sup>1</sup>Department of Aquaculture, Faculty of Fisheries, Pekalongan University, Pekalongan, Indonesia

<sup>2</sup>Department of Aquaculture, Faculty of Agriculture and Natural Resources, An Giang University, Long Xuyen City, Vietnam

<sup>7</sup>Corresponding author: [ariadi\\_heri@yahoo.com](mailto:ariadi_heri@yahoo.com)

Received: 26 August 2023

Accepted: 9 November 2023

## ABSTRACT

*Chlamydomonas* spp and *Chlorella* spp are plankton genera that can be used to determine the level of balance in the aquatic environment. the purpose of this study is to determine the abundance of plankton *Chlamydomonas* spp and *Chlorella* spp and its relation to the clinical performance of red tilapia in silvofishery ponds. The research method used is an ex-*post* *facto* causal design with random sampling. The results showed that the water quality conditions in the silvofishery ponds were very good except for the Nitrate parameters of 0.00-1.50 mg/L and Salinity of 2-11 gr/L which exceeded the quality standards. In the silvofishery ponds, 5 plankton classes were found, namely Chlorophyceae, Cyanophyceae, Chrysophyceae, Protozoa, and Dinophysis, consisting of 15 genera. The most dominant class of Chlorophyceae with an abundance of 2.88E+06 cells/ml. The Chlorophyceae class is dominated by *Chlorella* spp. (2.63E+06 cell/ml) and *Chlamydomonas* spp. (2.20E+05 cells/ml). The abundance of *Chlamydomonas* spp. in silvofishery ponds is closely related to phosphate solubility (0.988). Based on observations of clinical symptoms in the eyes, gills, and fins of fish, no disease infections or physical abnormalities were found in fish cultured in silvofishery ponds. Fish tend to live well with a growth rate of 0.44 gr/day. The aquatic ecosystem of the silvofishery pond which is dominated by the plankton *Chlorella* spp. and *Chlamydomonas* spp had a good effect

on the performance and condition of the fish being cultivated. This study concludes that the abundance of *Chlamydomonas* spp and *Chlorella* spp was found to be very dominant compared to other plankton genera. The presence of *Chlamydomonas* spp and *Chlorella* spp illustrates good and stable environmental conditions in silvofishery ponds followed by no clinical signs of disease infection in fish reared during the study period..

**KEYWORDS:** Chlorophyceae; silvofishery; *O. niloticus*; plankton; symptom.

## INTRODUCTION

Silvofishery is a multi-organism aquaculture model in mangrove forest ecosystems (Bao et al, 2013; Herrera et al, 2015). The advantages of silvofishery include better harvest productivity levels, low aquaculture waste, and minimal aquaculture inputs (Musa et al, 2020). The concept of silvofishery can be developed in various types of waters with various commodities (Lukman et al, 2021). The silvofishery concept is very suitable to be developed in tropical waters (Ariadi et al, 2019).

*Chlamydomonas* spp and *Chlorella* spp are types of plankton that are widely used for aquaculture activities (Che and Kim, 2023). *Chlamydomonas* spp and *Chlorella* spp are microalgae that can be used as bio-indicators of the aquatic environment (Zhou et al, 2022). *Chlamydomonas* spp is a multicellular microalga that can live in extreme waters (Zhang et al, 2021). *Chlorella* spp is a plankton that grows continuously in tropical waters (Hernandez et al, 2006; Alagawany et al, 2021).

The existence of *Chlamydomonas* sp and *Chlorella* spp is very important as bio-indicator of the aquatic environment (Khalil et al, 2021; Zhou et al, 2022). A good aquatic environment will affect the productivity level of aquaculture (Dong et al, 2022). Conversely, a bad aquatic environment will make fish susceptible to stress and die (Britton et al, 2023). One way to detect the quality level of a water site is to see the abundance of plankton such as *Chlamydomonas* spp and *Chlorella* spp (Soeprapto et al, 2023).

Red tilapia is a type of fish that is widely aquaculture in tropical waters (Ekasari et al, 2023; Arias et al, 2023). Red tilapia is a type of fish that is reactive to changes in aquatic environmental conditions (Nguyen et al, 2023). One indicator of the response of red tilapia to environmental conditions can be seen in the clinical symptoms in the physical body (Wang et al, 2022). The presence of clinical symptoms in the tilapia body indicates a physiological reaction as an adaptation to a changing environmental habitat (Wang et al, 2022; Lukman et al, 2023). Clinical symptoms on the body of red tilapia can be seen from the presence of spots,

changes in body structure, color changes, and the presence of wounds (Arias et al, 2022). From this clinical response, we can describe the condition of the red tilapia habitat waters (Ariadi and Abidin, 2019).

It is important to understand the existence of *Chlamydomonas* spp and *Chlorella* spp as environmental bio-indicators and clinical symptoms of fish as a physiological response in pathology, so the purpose of this study is to determine the abundance of plankton *Chlamydomonas* spp and *Chlorella* spp and its relation to the clinical performance of red tilapia in ponds silvofishery. From the results of this study, we can determine whether there is a correlation between the presence of aquatic bio-indicator microorganisms on clinical symptoms in the bodies of aquaculture in silvofishery ponds.

## 1 MATERIALS AND METHODS

This research was conducted in the silvofishery pond of Slamaran Village, Pekalongan City with the *ex-pose facto causal* design concept or direct data collection in the field without any treatment engineering. Sampling of fish, water quality, plankton, and fish clinical symptoms was carried out by random sampling. Parameters observed were clinical symptoms on the physical body of the fish, abundance of plankton, and water quality in silvofishery ponds.

The parameters of the clinical symptoms observed were changes in the physical structure of the fish, body deformities, injuries, and changes in the color of the fish. Fish samples were observed by taking fish populations floating on the surface of the water. The water quality parameters observed were pH, dissolved oxygen, salinity, temperature, phosphate, nitrite, nitrate, and ammonia. The plankton parameters observed were the plankton abundance index which was calculated using the Olympus CX23 microscope and the NEUBEUER© Haemocytometer which was then calculated using the formula by APHA, (1998) as follows:

$$N = Z \times \frac{x}{y} \times \frac{1}{V}$$

Information :

N : Plankton individually abundant (ind/ltr)

Z : Plankton individual number

X : Volume of water sample (40 mL)

Y : Volume of 1 water drop (0.06 mL)

V : Volume of filtered water (100 L)

Furthermore, the data is grouped based on the parameters and research indicators to test the data analysis. Data analysis was carried out descriptively using SPSS 1.16 software.

## RESULT AND DISCUSSION

### Water quality parameters

The water quality parameters observed in this study can be seen in Table 1. Based on a comparison of the measurement results with standard aquaculture water quality standards, it was found that only Nitrate parameters were 0.00-1.50 mg/L and Salinity 2-11 gr/L which did not meet the quality standard threshold values water for aquaculture activities. This means that the water conditions in silvofishery ponds tend to be low saline and eutrophic. The presence of high organic matter and low salinity levels makes aquatic ecosystems tend to be eutrophic (Zhang et al, 2023).

Water quality in silvofishery ponds tends to be better and more stable than conventional ponds. The existence of a mangrove ecosystem makes the waste absorption process more intense (Musa et al, 2020). The mangrove ecosystem functions as a biofilter and catcher of waste inorganic compounds (Cunha et al, 2019). The mangrove ecosystem in silvofishery cultivation also functions as a natural habitat for crabs and fish roaming places (Sarower et al, 2021).

Table 1. Water quality parameters in silvofishery ponds

Parameter	Measurement Results	Standard*
Phosphate	0.03-0.10	<0.10
Ammonia	0.01-0.15	<0.10
Nitrite	0.00-0.10	<1.00
Nitrate	0.50-1.50	<1.00
Temperature	26.25-31.40	24-32
Salinity	2-11	25-35
Dissolved Oxygen	4.50-20.00	>4
pH	7.5-8.2	7.5-8.5

\*Ariadi et al, (2021)

4 The abundance of plankton in silvofishery ponds can be seen in Table 2. Based on the research data, 5 different classes and 15 genera of plankton were found. The most diverse plankton are from the class Dinophyceae in which 5 genera were found (Table 2.). The diversity of Dinophysis is quite diverse because the sampling was carried out during the dry season (Ajani et al, 2022). The genus Dinophyceae found in silvofishery ponds are *Chryptomonas* spp., *Gymnodinium* spp., *Noctiluca* spp., *Peridinium* spp., *Prorocentrum* spp.

In the silvofishery ponds, the plankton genera Chlorophyceae, Cyanophyceae, Chrysophyceae, and Protozoa were also found. This means that the level of plankton diversity in silvofishery ponds is quite high. The existence of a balance of chemical, physical and biological parameters greatly supports the stability of plankton diversity in the waters (Geng et al, 2022). Plankton will experience seasonal succession which allows the dominance of certain classes (David et al, 2020; Ajani et al, 2022).


Table 2. Class and genus of plankton in silvofishery ponds

Classess	Genus
Chlorophyceae	<i>Dicthyospaerium</i> sp., <i>Chlamydomonas</i> sp., <i>Chlorella</i> sp.
Cyanophyceae	1 <i>Microcystis</i> sp., <i>Oscillatoria</i> sp.
Chrysophyceae	<i>Amphora</i> sp., <i>Diploneis</i> sp.
Dinophyceae	<i>Chryptomonas</i> sp., <i>Gymnodinium</i> sp., <i>Noctiluca</i> sp., <i>Peridinium</i> sp., <i>Prorocentrum</i> sp.
Protozoa	<i>Acantocystis</i> sp., <i>Ciliata</i> sp., <i>Euplotes</i> sp.

### Plankton Classes

In silvofishery ponds, 5 classes of plankton were found, namely Chlorophyceae, Cyanophyceae, Chrysophyceae, Dinophyceae, and Protozoa (Figure 1.). The existence of plankton in aquatic ecosystems is determined by the abundance of nutrients (Soeprapto et al, 2023). The Chlorophyceae class indicates that silvofishery pond waters are eutrophic. The level of the trophic status of the waters is determined by geographical conditions, wind patterns, and seasons (Valentin et al, 2021). The more fertile the waters, the higher the succession of plankton at the genus and species level (Leles et al, 2021).

Chlorophyceae live in colonies and their life pattern is determined by the level of temperature distribution (Ariadi et al, 2019). The high abundance of Chlorophyceae also allows for more intense dissolved oxygen production (Wafi et al, 2021). These conditions also have an impact on diurnal pH fluctuations (Soeprapto et al, 20.23). Chlorophyceae will carry out photosynthesis during the day and carry out quite intense respiration at night (Yinding et al, 2021).




**Figure 1.** Plankton abundance in silvofishery ponds by class

### *Chloropyceae*

Chlorophyceae in silvofishery ponds have the highest dominance level compared to other classes. The plankton genera in the Chlorophyceae class found included *Dicthyosphaerium* spp., *Chlamydomonas* spp., *Chlorella* spp. (Fig. 2.). Most of the plankton genera of the Chlorophyceae class have similar morphological and ecological characteristics (Fucikova and Lewis, 2016; Teemrалеva and Bukin, 2022). The abundance of Chlorophyceae in waters is influenced by temperature, chlorophyll-a, and light (Rocha et al, 2021).

*Chlorella* spp. is so dominant compared to *Dicthyosphaerium* spp, and *Chlamydomonas* spp. *Chlorella* spp is a cosmopolitan photosynthetic plankton genus (Gong et al, 2021). The life cycle of cosmopolitan plankton is largely determined by temperature distribution and the presence of nutrients as limiting factors (Ariadi et al, 2019; Ariadi et al, 2022). In aquaculture ponds, *Chlorella* spp can be used as a natural food by fish (Saldana et al, 2022).



**Figure 2.** The abundance of the plankton genus Chlorophyceae in silvofishery ponds

The high abundance of *Chlorella* spp. ( $2.63E+06$  cell/ml) and *Chlamydomonas* spp. ( $2.20E+05$  cell/ml) can be used as an environmental bio-indicator for aquaculture waters (Che and Kim, 2023). This means that the aquatic ecosystem in silvofishery ponds is very fertile. The high level of primary productivity is caused by the accumulation of aquaculture waste, organic matter, and the presence of trace mineral elements (Ray et al, 2023). Intensive pattern fish farming activities tend to have a eutrophic impact on the aquatic environment of ponds (Han and Chui, 2016).

#### *Chlorella* spp and *Chlamydomonas* spp


*Chlorella* spp. and *Chlamydomonas* spp. are a genus of Chlorophyceae that often grows in aquaculture waters (Ariadi et al, 2021). The abundance of *Chlorella* spp. and *Chlamydomonas* spp. in silvofishery ponds is presented in Figure 3. The abundance of *Chlorella* spp. ranged from  $2.80E+05$  -  $8.70E+05$  cell/ml and *Chlamydomonas* spp.  $2.00E+04$  -  $9.00E+04$  cells/ml. Increasing the biomass of *Chlorella* spp and *Chlamydomonas* spp. caused by a decrease in water salinity levels and an increase in extracellular polymeric substance (EPS) compounds (Vo et al, 2020).

The presence of *Chlorella* spp. and *Chlamydomonas* spp. indicates a low level of solubility of organic matter in aquatic ecosystems (Bilal et al, 2019). *Chlorella* spp. live well in heterotrophic waters (Wang et al, 2016). *Chlamydomonas* spp. is a type of algae that can effectively absorb  $NH_4$  in aquatic ecosystems (Zhou et al, 2022). This means that the presence



of the two plankton genera indicates a stable silvofishery aquatic ecosystem with moderate nutrient abundance.

The level of abundance ratio between *Chlorella* spp. and *Chlamydomonas* spp are 12:1 or *Chlorella* spp so it exists in silvofishery aquatic ecosystems. Differences in life cycles and the effectiveness of nutrient utilization are the differentiating factors for plankton growth in the waters (Cao et al, 2021; Zhou et al, 2022). *Chlamydomonas* spp. is a type of plankton that can grow in phototrophic, heterotrophic, and mixotrophic waters (Moon et al, 2013). These conditions allow *Chlamydomonas* spp to grow in various water conditions. Likewise, *Chlorella* spp is solitary and its growth is influenced by the level of temperature distribution (Ariadi et al, 2019; Serra-Maia et al, 2016).



**Figure 3.** The abundance of *Chlamydomonas* spp. and *Chlorella* spp. in silvofishery ponds

Correlation test results between the abundance of *Chlorella* spp. and *Chlamydomonas* spp. on water nutrient parameters (NH<sub>3</sub>, PO<sub>4</sub>, NO<sub>2</sub>, NO<sub>3</sub>) showed only the phosphate parameter (0.988) which correlated with the abundance of *Chlamydomonas* spp. (Table 2.). For other parameters, there is no strong correlation. The increase in element P will make the metabolic system of *Chlamydomonas* spp in carrying out assimilation more intense (Plouviez et al, 2023). Excessive abundance of phosphate will damage *Chlorella* spp cells but in *Chlamydomonas* spp phosphate becomes a growth-limiting factor (Shilton et al, 2012; Li et al, 2018). The presence of phosphate as a limiting element will be very important for aquatic ecosystems (Ray et al,

2023). Phosphate will dissolve easily and will precipitate if the lime content in the water is relatively high.

Table 1. Correlation test results between the abundance of *Chlorella* spp and *Chlamydomonas* spp on water nutrient parameters

		<i>Chlamydomonas</i>	<i>Chlorella</i>	NH <sub>3</sub>	PO <sub>4</sub>	NO <sub>2</sub>	NO <sub>3</sub>
		spp	spp.				
<i>Chlamydomonas</i> spp	Pearson	1	.717	.900	.988*	.256	.473
	Correlation		.283	.100	.012	.744	.527
	Sig.(2-tailed)	4	4	4	4	4	4
	N						
<i>Chlorella</i> spp.	Pearson	.717	1	.944	.646	.205	.802
	Correlation	.283		.056	.354	.795	.198
	Sig.(2-tailed)	4	4	4	4	4	4
	N						

Overall, with good water quality parameters in silvofishery ponds, it was found that the highest level of plankton dominance came from the Chlorophyceae class. *Chlorella* spp. and *Chlamydomonas* spp. is the dominant genus in the class Chlorophyceae in silvofishery ponds. Chlorophyceae are cosmopolitan in nutrient-rich environments (Ariadi et al, 2023). The high load of feed waste and the accumulation of nutrients will trigger the growth of *Chlorella* spp. biomass. and *Chlamydomonas* spp. in aquatic ecosystems (Xie et al, 2019; Ullmann and Grimm, 2021). This is consistent with the results of the correlation test showing a high closeness relationship between the abundance of *Chlamydomonas* spp and the solubility of phosphate in waters. Another factor that causes *Chlorella* spp. and *Chlamydomonas* spp. dominant is the temperature level and the value of the salinity of the waters (Ma et al, 2020).

#### Clinical Symptoms and Fish Growth

Clinical symptoms of red tilapia reared in silvofishery ponds can be seen in Figure 4. Physically the fish do not experience any symptoms or structural changes in their physical organs. The color of the gills still looks bright red (4a), the color of the eyes is still bright (4b), and the dorsal fin is intact and has not undergone any structural changes (4c). This means that

fish kept in silvofishery ponds are healthy and do not experience clinical symptoms of changes in the structure of their organs. Clinical symptoms in fish are usually caused by infection with pathogenic bacteria and abnormal changes in behavior (Okon et al, 2023; Amminger et al, 2023). plankton and pathogenic bacteria have complementary properties in the host infection process (Costello et al, 2023).

There are no bad clinical symptoms in fish due to good environmental habitat conditions (Mramba and Kahindi, 2023). Good environmental conditions can be seen from the condition of water quality and the level of abundance of plankton that grows. Plankton and water quality are natural indicators that can stabilize aquaculture ecosystems (Meng et al, 2022). The abundance of *Chlorella* spp. and *Chlamydomonas* spp. which is a genus of beneficial plankton will affect the performance of fish cultivated. *Chlorella* spp. and *Chlamydomonas* spp. is a good aquatic environment bio-indicator, so there is a correlation that good environmental conditions have an impact on the physical performance of fresher fish (Ariadi et al, 2020).

No clinical symptoms were found on the body of the fish indicating that the silvofishery concept is very feasible to be developed. The indicators are the lack of symptoms of physical defects in the body of the fish being cultivated and the relatively good condition of the water quality in silvofishery ponds (Musa et al, 2020). In addition, the growth of profitable plankton genera greatly supports the productivity of the operational cycle of silvofishery. *Chlorella* spp and *Chlamydomonas* spp are types of plankton that are widely used as natural feed and aquatic bio-indicator agents. (Irihimovitch et al, 2008).




**Figure 4.** Clinical symptoms on the fish body a.) gills, b.) eyes, c.) dorsal fin

Data from observations of fish growth during the first 40 days of the cultivation period can be seen in Figure 5. Fish growth rates ranged from 3.21-20.76 g and continued to increase (Figure 5.). The growth of the biomass of aquaculture organisms will continue to increase as the cultivation period increases (Ariadi et al, 2019). Fish growth factors are influenced by the

effectiveness of feeding, feed quality, and supportive environmental habitat conditions (Pouil et al, 2023).

The average growth rate of fish is 0.44 gr/day. This figure is quite high for the development of silvofishery. The average growth rate of fish is a visualization of the effectiveness of feed management given during the period (Lee et al, 2023). Effective feeding and good feed quality play an important role in fish growth performance (Madusari et al, 2022). Thus, it can be said that a silvofishery system with a good fish-rearing model will have a progressive increase in fish biomass and minimal impact on environmental pollution. The minimal impact of waste on the environment will provide a higher carrying capacity rate (Ariadi et al, 2023)



**Figure 5.** Growth rate of fish in silvofishery ponds during the study

The results of the research study as a whole show that the conditions of water quality and the presence of plankton in silvofishery ponds are ecologically quite good. In addition, the presence of plankton *Chlorella* spp. and *Chlamydomonas* spp. Dominant ones are considered good enough to support aquaculture activities. The concept of silvofishery with minimal waste pollution affects the stability of the ecosystem in pond waters (Wijaya et al, 2019). These good ecological conditions affect the performance of the fish being cultivated (Ariadi, 2023). Aquaculture in silvofishery ponds is very healthy and there are no clinical symptoms due to disease infection. The more stable ecosystem conditions in silvofishery ponds provide a reduction in the prevalence rate of disease spread (De-Leon-Herrera et al, 2015). In addition, the growth rate of aquaculture in silvofishery ponds tends to increase its biomass progressively every day. This means that the fish cultivated in this silvofishery pond are very good and healthy so they are very worthy of being developed (Mirera, 2011). The healthy condition of

fish can be seen from the lack of clinical symptoms due to diseases acquired during the cultivation period.

## CONCLUSION

The abundance of *Chlamydomonas* spp and *Chlorella* spp was found to be very dominant compared to other plankton genera. The presence of *Chlamydomonas* spp and *Chlorella* spp illustrates good and stable environmental conditions in silvofishery ponds followed by no clinical signs of disease infection in fish reared during the study period.

## ACKNOWLEDGMENT

The author would like to thank the Adaptation Programme with Kemitraan Partnership Foundation for facilitating this research through contract number PSC-003/2023.

## LITERATUR CITED

- Ajani, P.A., Henriquez-Nunez, H.F., Verma, A., Nagai, S., Uchida, H., Tesoriero, M.J., Farrell, H., Zammit, A., Brett, S., Murray, S.A. (2022). Mapping the development of a Dinophysis bloom in a shellfish aquaculture area using a novel molecular qPCR assay. **Harmful Algae**, 116, 102253. <https://doi.org/10.1016/j.hal.2022.102253>
- Alagawany, M., Taha, A.E., Noreldin, A., El-Tarabily, K.A., El-Hack, M.E.A. (2021). Nutritional applications of species of Spirulina and Chlorella in farmed fish: A review. **Aquaculture**, 542, 736841. <https://doi.org/10.1016/j.aquaculture.2021.736841>
- Amminger, G.P., Rice, S., Davey, C.G., Quinn, A.L., Hermens, D.F., Zmicerevska, N., Nichles, A., Hickie, I., Incerti, L., Weller, A., Joseph, S., Hilton, Z., Pugh, C., Rayner, M., Reid, N., Ratheesh, A., Yung, A.R., Yuen, H.P., Mackinnon, A., Hetrick, S., Lin, A. (2023). The Addition of Fish Oil to Cognitive Behavioural Case Management for Youth Depression (YoDA-F): A Randomised, Double-Blind, Placebo-Controlled, Multicentre Clinical Trial. **Biological Psychiatry**, 1-12. <https://doi.org/10.1016/j.biopsych.2023.06.015>
- Ariadi, H., Azril, M., Mujtahidah, T. (2023). Water Quality Fluctuations In Whiteleg Shrimp (*Litopenaeus vannamei*) Cultivation During The Dry And Rainy Seasons. **Croatian Journal of Fisheries: Ribarstvo**, 81(3), 127-137.
- Ariadi, H. (2023). **Dinamika Wilayah Pesisir**. Malang: UB Press. 157 pp. (in indonesian)
- <sup>3</sup> Ariadi, H., and Abidin, Z. (2019). Study of partnership pattern among farmers of tilapia fish (*Oreochromis niloticus*) and fish breeding Centre Klemunan in Wlingi of Blitar

- Regency. **ECOSOFIM: Economic and Social of Fisheries and Marine Journal**, 6(02), 194-201.
- Ariadi, H., Mahmudi, M., Fadjar, M. (2019). Correlation between density of vibrio bacteria with *Oscillatoria* sp. abundance on intensive *Litopenaeus vannamei* shrimp ponds. **Research Journal of Life Science**, 6(2), 114-129.
- Ariadi, H., Fadjar, M., Mahmudi, M. (2019). Financial feasibility analysis of shrimp vannamei (*Litopenaeus vannamei*) culture in intensive aquaculture system with low salinity. **ECOSOFiM (Economic and Social of Fisheries and Marine Journal)**, 7(01), 95-108.
- Ariadi, H., Fadjar, M., Mahmudi, M. (2019). The relationships between water quality parameters and the growth rate of white shrimp (*Litopenaeus vannamei*) in intensive ponds. **Aquaculture, Aquarium, Conservation & Legislation**, 12(6), 2103-2116.
- Ariadi, H., Pandaingan, I.A.H., Soeprijanto, A., Maemunah, Y., Wafi, A. (2020). Effectiveness of using Pakcoy (*Brassica rapa* L.) and Kailan (*Brassica oleracea*) plants as vegetable media for aquaponic culture of Tilapia (*Oreochromis* sp.). **Journal of Aquaculture Development and Environment**, 3(2), 156-162. DOI: <http://dx.doi.org/10.31002/jade.v3i2.3320>
- Ariadi, H., Wafi, A., Madusari, B.D. (2021). **Dynamics of Dissolved Oxygen (Case Study on Shrimp Cultivation)**. ADAB Corp. Indramayu. 138 pp. (in indonesian)
- Ariadi, H., Syakirin, M.B., Hidayati, S., Madusari, B.D., Soeprapto, H. (2022). Fluctuation Effect of Dissolved of TAN (Total Ammonia Nitrogen) on Diatom Abundance in Intensive Shrimp Culture Ponds. **IOP Conference Series: Earth and Environmental Science**, 1118(1), 012001. DOI 10.1088/1755-1315/1118/1/012001
- Arias, L., Marquez, D.M., Zapata, J.E. (2022). Quality of red tilapia viscera oil (*Oreochromis* sp.) as a function of extraction methods. **Heliyon**, 8(5), e09546. <https://doi.org/10.1016/j.heliyon.2022.e09546>
- Arias, L., Sanchez-Henao, C.P., Zapata, J.E. (2023). Optimization of the separation conditions of antioxidant peptides from red tilapia (*Oreochromis* spp.) viscera on cross-flow filtration ceramic membranes. **South African Journal of Chemical Engineering**, 45, 100-110. <https://doi.org/10.1016/j.sajce.2023.05.002>
- Bao, H., Wu, Y., Unger, D., Du, J., Herbeck, L.S., Zhang, J. (2013). Impact of the conversion of mangroves into aquaculture ponds on the sedimentary organic matter composition in a tidal flat estuary (Hainan Island, China). **Continental Shelf Research**, 57, 82-91. <https://doi.org/10.1016/j.csr.2012.06.016>

- Bilal, M., Adeel, M., Rasheed, T., Zhao, Y., Iqbal, H.M.N. (2019). Emerging contaminants of high concern and their enzyme-assisted biodegradation – A review. **Environment International**, *124*, 336-353. <https://doi.org/10.1016/j.envint.2019.01.011>
- Britton, J.R., Andreou, D., Lopez-Bejar, M., Carbajal, A. (2023). Relationships of scale cortisol content suggest stress resilience in freshwater fish vulnerable to catch-and-release angling in recreational fisheries. **Fisheries Research**, *266*, 106776. <https://doi.org/10.1016/j.fishres.2023.106776>
- Cao, Y., Shao, P., Chen, Y., Zhou, X., Yang, L., Shi, H., Yu, K., Luo, X., Luo, X. (2021). A critical review of the recovery of rare earth elements from wastewater by algae for resources recycling technologies. **Resources, Conservation and Recycling**, *169*, 105519. <https://doi.org/10.1016/j.resconrec.2021.105519>
- Che, C.A., and Kim, S.K. (2023). The optimization of culture conditions for the enhancement of biomass and lipids in *Chlamydomonas hedleyi* using a two-phase culture system. **Bioresource Technology Reports**, *21*, 101342. DOI: 10.1016/j.biteb.2023.101342
- Costello, K.E., Haberlin, D., Lynch, S.A., McAllen, R., O’Riordan, R.M., Culloty, S.C. (2023). Regional differences in zooplankton-associated bacterial communities and aquaculture pathogens across two shelf seas. **Estuarine, Coastal and Shelf Science**, *281*, 108179. <https://doi.org/10.1016/j.ecss.2022.108179>
- Cunha, M.E., Quental-Ferreira, H., Parejo, A., Gamito, S., Ribeiro, L., Moreira, M., Monteiro, I., Soares, F., Pousao-Ferreira, P. (2019). Understanding the individual role of fish, oyster, phytoplankton and macroalgae in the ecology of integrated production in earthen ponds. **Aquaculture**, *512*, 734297. <https://doi.org/10.1016/j.aquaculture.2019.734297>
- David, V., Tortajada, S., Savoye, N., Breret, M., Lachaussee, N., Philippine, O., Robin, F.X., Dupuy, C. (2020). Impact of human activities on the spatio-seasonal dynamics of plankton diversity in drained marshes and consequences on eutrophication. **Water Research**, *170*, 115287. <https://doi.org/10.1016/j.watres.2019.115287>
- De-Leon-Herrera, R., Flores-Verdugo, F., Flores-de-Santiago, F., Gonzalez-Farias, F. (2015). Nutrient removal in a closed silvofishery system using three mangrove species (*Avicennia germinans*, *Laguncularia racemosa*, and *Rhizophora mangle*). **Marine Pollution Bulletin**, *91(1)*, 243-248. <https://doi.org/10.1016/j.marpolbul.2014.11.040>
- Dong, S., Shan, H., Yu, L., Liu, X., Ren, Z., Wang, F. (2022). An ecosystem approach for integrated pond aquaculture practice: Application of food web models and ecosystem

- indices. **Ecological Indicators**, *141*, 109154.  
<https://doi.org/10.1016/j.ecolind.2022.109154>
- Ekasari, J., Napitupulu, A.D., Djurstedt, M., Wiyoto, W., Baruah, K., Kiessling, A. (2023). Production performance, fillet quality and cost effectiveness of red Tilapia (*Oreochromis* sp.) culture in different biofloc systems. **Aquaculture**, *563*, 738956.  
<https://doi.org/10.1016/j.aquaculture.2022.738956>
- Fucikova, K., and Lewis, L.A., Lewis, P.O. (2016). Comparative analyses of chloroplast genome data representing nine green algae in Sphaeropleales (Chlorophyceae, Chlorophyta). **Data in Brief**, *7*, 558-570. <https://doi.org/10.1016/j.dib.2016.03.014>
- Geng, Y., Li, M., Yu, R., Sun, H., Zhang, L., Sun, L., Lv, C., Xu, J. (2022). Response of planktonic diversity and stability to environmental drivers in a shallow eutrophic lake. **Ecological Indicators**, *144*, 109560. <https://doi.org/10.1016/j.ecolind.2022.109560>
- Gong, Y., Zheng, X., Huang, J. (2021). *Chlorella* sp. Mg shows special trophic transitions and biomass production. **Bioresource Technology Reports**, *16*, 100854.  
<https://doi.org/10.1016/j.biteb.2021.100854>
- Han, Z., and Chui, B. (2016). Performance of macrophyte indicators to eutrophication pressure in ponds. **Ecological Engineering**, *96*, 8-19.  
<https://doi.org/10.1016/j.ecoleng.2015.10.019>
- Hernandez, J.P., de Bashan, L.E., Bashan, Y. (2006). Starvation enhances phosphorus removal from wastewater by the microalga *Chlorella* spp. co-immobilized with *Azospirillum brasilense*. **Enzyme and Microbial Technology**, *38*, 190-198.  
<https://doi.org/10.1016/j.enzmictec.2005.06.005>
- Herrera, R.D.L., Flores-Verdugo, F., de Santiago, F.F., Farias, F.G. (2015). Nutrient removal in a closed silvofishery system using three mangrove species (*Avicennia germinans*, *Laguncularia racemosa*, and *Rhizophora mangle*). **Marine Pollution Bulletin**, *91(1)*, 243-248. <https://doi.org/10.1016/j.marpolbul.2014.11.040>
- Irihimovitch, V., Yehudai, S., Resheff. (2008). Phosphate and sulfur limitation responses in the chloroplast of *Chlamydomonas reinhardtii*. **FEMS Microbiology Letters**, *283*, 1-8. <https://doi.org/10.1111/j.1574-6968.2008.01154.x>
- Khalil, S., Mahnashi, M.H., Hussain, M., Zafar, N., Nisa, W.U., Khan, F.S., Afzal, U., Shah, G.M., Niazi, U.M., Awais, M., Irfan, M. (2021). Exploration and determination of algal role as Bioindicator to evaluate water quality – Probing fresh water algae. **Saudi Journal of Biological Sciences**, *28(10)*, 5728-5737.  
<https://doi.org/10.1016/j.sjbs.2021.06.004>



- 5  
Lee, M.J., Kim, J., Baek, S.I., Cho, S.H. (2023). Substitution effect of fish meal with meat meal in diet on growth performance, feed consumption, feed utilization, chemical composition, hematology, and innate immune responses of rockfish (*Sebastes schlegeli*). **Aquaculture**, 571, 739467. <https://doi.org/10.1016/j.aquaculture.2023.739467>
- Leles, S.G., Bruggeman, J., Polimene, L., Blacford, J., Flynn, K.J., Mitra, A. (2021). Differences in physiology explain succession of mixoplankton functional types and affect carbon fluxes in temperate seas. **Progress in Oceanography**, 190, 102481. <https://doi.org/10.1016/j.pocean.2020.102481>
- Li, Q., Fu, L., Wang, Y., Zhou, D., Rittmann, B.E. (2018). Excessive phosphorus caused inhibition and cell damage during heterotrophic growth of *Chlorella regularis*. **Bioresource Technology**, 268, 266-270. <https://doi.org/10.1016/j.biortech.2018.07.148>
- Lukman, K.M., Uchiyama, Y., Kohsaka, R. (2021). Sustainable aquaculture to ensure coexistence: Perceptions of aquaculture farmers in East Kalimantan, Indonesia. **Ocean & Coastal Management**, 213, 105839.
- Lukman, B., Roslindawani, M.N., Azzam-Sayuti, M., Norfarrah, M.A., Annas, S., Ina-Salwany, M.Y., Zamri-Saad, M., Nor-Yasmin, A.R., Amin-Nordin, S., Barkham, T., Amal, M.N.A. (2023). Disease development in red hybrid tilapia following single and co-infection with tilapia lake virus and *Streptococcus agalactiae*. **Aquaculture**, 567, 739251. <https://doi.org/10.1016/j.aquaculture.2023.739251>
- Ma, R., Zhao, X., Ho, S.H., Shi, X., Liu, L., Xie, Y., Chen, J., Lu, Y. (2020). Co-production of lutein and fatty acid in microalga *Chlamydomonas* sp. JSC4 in response to different temperatures with gene expression profiles. **Algal Research**, 47, 101821. <https://doi.org/10.1016/j.algal.2020.101821>
- Madusari, B.D., Ariadi, H., Mardhiyana, D. (2022). Effect of the feeding rate practice on the white shrimp (*Litopenaeus vannamei*) cultivation activities. **Aquaculture, Aquarium, Conservation & Legislation-International Journal of the Bioflux Society**, 15(1), 473-479.
- Meng, C., Liu, H., Li, Y., Shen, J., Li, X., Wu, J. (2022). Effects of environmental and agronomic factors on pond water quality within an intensive agricultural landscape in subtropical southern China. **Agricultural Water Management**, 274, 107953. <https://doi.org/10.1016/j.agwat.2022.107953>

- Mirera, O.D. (2011). Trends in exploitation, development and management of artisanal mud crab (*Scylla serrata*-Forsskal-1775) fishery and small-scale culture in Kenya: An overview. **Ocean & Coastal Management**, 54(11), 844-855. <https://doi.org/10.1016/j.ocecoaman.2011.08.001>
- Moon, M., Kim, C.W., Park, W.K., Yoo, G., Choi, Y.E., Yang, J.W. (2013). Mixotrophic growth with acetate or volatile fatty acids maximizes growth and lipid production in *Chlamydomonas reinhardtii*. **Algal Research**, 2(4), 352-357. <https://doi.org/10.1016/j.algal.2013.09.003>
- Mramba, R.P., and Kahindi, E.J. (2023). Pond water quality and its relation to fish yield and disease occurrence in small-scale aquaculture in arid areas. **Heliyon**, 9, e16753. <https://doi.org/10.1016/j.heliyon.2023.e16753>
- Musa, M., Mahmudi, M., Arsad, S., Buwono, N.R. (2020). Feasibility study and potential of pond as silvofishery in coastal area: Local case study in Situbondo Indonesia. **Regional Studies in Marine Science**, 33, 100971. <https://doi.org/10.1016/j.rsma.2019.100971>
- Nguyen, M.T., Pham, N.T.A., Vo, L.T., Truong, D.V., Nguyen, H.V., Nguyen, T.D.Q., Nguyen, P.N., Bossier, P. (2023). Integrated mariculture of co-cultured whiteleg shrimp (*Litopenaeus vannamei*) and grey mullet (*Mugil cephalus*) in sequence with red tilapia (*Oreochromis* spp.) in a closed biofloc-based system. **Aquaculture**, 566, 739200. <https://doi.org/10.1016/j.aquaculture.2022.739200>
- Okon, E.M., Okocha, R.C., Taiwo, A.B., Michael, F.B., Bolanle, A.M. (2023). Dynamics of co-infection in fish: A review of pathogen-host interaction and clinical outcome. **Fish and Shellfish Immunology Reports**, 4, 100096. <https://doi.org/10.1016/j.fsirep.2023.100096>
- Plouviez, M., Bolot, P., Shilton, A., Guieysse, B. (2023). Phosphorus uptake and accumulation in *Chlamydomonas reinhardtii*: Influence of biomass concentration, phosphate concentration, phosphorus depletion time, and light supply. **Algal Research**, 71, 103085. <https://doi.org/10.1016/j.algal.2023.103085>
- Pouil, S., Kerneis, T., Quillet, E., Labbe, L., Lallias, D., Phocas, F., Nivet, M.D. (2023). Isogenic lines of rainbow trout (*Oncorhynchus mykiss*) as a tool to assess how growth and feeding behaviour are correlated to feed efficiency in fish. **Aquaculture**, 577, 739904. <https://doi.org/10.1016/j.aquaculture.2023.739904>
- Ray, L.I.P., Swetha, K., Singh, A.K., Singh, N.J. (2023). Water productivity of major pulses – A review. **Agricultural Water Management**, 281, 108249. <https://doi.org/10.1016/j.agwat.2023.108249>

- Ray, A., Kumar, M., Karim, A.A., Biswas, K., Mohanty, S., Shadangi, K.P., Kumar, S., Sarkar, B. (2023). Potassium-phosphorus-sulfur augmented biochar production from potentially toxic elements abated gypsum pond wastewater of phosphate fertilizer industry. **Journal of Environmental Chemical Engineering**, *11(5)*, 110404. <https://doi.org/10.1016/j.jece.2023.110404>
- Rocha, G.S., Parrish, C.C., Espindola, E.L.G. (2021). Effects of copper on photosynthetic and physiological parameters of a freshwater microalga (Chlorophyceae). **Algal Research**, *54*, 102223. <https://doi.org/10.1016/j.algal.2021.102223>
- Saldana, K., Angulo, E., Mercado, I., Castellar, G., Cubillan, N. (2022). Removal of minocycline from high concentrated aqueous medium by nonliving and lipid-free *Chlorella* sp. biomass. **Bioresource Technology Reports**, *17*, 100921. <https://doi.org/10.1016/j.biteb.2021.100921>
- Sarower, M.G., Al-Hasan, M.M., Rahman, M.S., Hasan, M.M., Ahmmed, M.K., Ali, M.Y., Giteru, S.G., Banu, G.R. (2021). Comparative growth and morphometric assessment between cultures of wild and hatchery-produced mud crabs. **Heliyon**, *7(9)*, e07964. <https://doi.org/10.1016/j.heliyon.2021.e07964>
- Serra-Maia, R., Bernard, O., Goncalves, A., Bensalem, S., Lopes, F. (2016). Influence of temperature on *Chlorella vulgaris* growth and mortality rates in a photobioreactor. **Algal Research**, *18*, 352-359. <https://doi.org/10.1016/j.algal.2016.06.016>
- Shilton, A.N., Powell, N., Guieysse, B. (2012). Plant based phosphorus recovery from wastewater via algae and macrophytes. **Current Opinion in Biotechnology**, *23(6)*, 884-889. <https://doi.org/10.1016/j.copbio.2012.07.002>
- Soeprapto, H., Ariadi, H., Badrudin, U. (2023). The abundance of *Microcystis* sp. on intensive shrimp ponds. **Depik**, *12 (1)*, , 105-110. DOI: 10.13170/depik.12.1.30433
- Soeprapto, H., Ariadi, H., Badrudin, U. (2023). The dynamics of *Chlorella* spp. abundance and its relationship with water quality parameters in intensive shrimp ponds. **Biodiversitas**, *24(5)*, 2919-2926. DOI <https://doi.org/10.13057/biodiv/d240547>
- Teemraleva, A.D., and Bukin, Y.S. (2022). Morphology, molecular phylogeny, and species delimitation within microalgal genera *Eubrownia*, *Spongiococcum*, and *Chlorococcum* (Chlorophyceae, Chlorophyta). **South African Journal of Botany**, *151*, 396-409. <https://doi.org/10.1016/j.sajb.2022.10.030>
- Ullmann, J., and Grimm, D. (2021). Algae and their potential for a future bioeconomy, landless food production, and the socio-economic impact of an algae industry. **Organic Agriculture**, *11*, 261–267.

- Valentin, J.L., Leles, S.G., Tenenbaum, D.R., Figueiredo, G.M. (2021). Frequent upwelling intrusions and rainfall events drive shifts in plankton community in a highly eutrophic estuary. **Estuarine, Coastal and Shelf Science**, 257, 107387. <https://doi.org/10.1016/j.ecss.2021.107387>.
- Vo, H.N.P., Ngo, H.H., Guo, W., Liu, Y., Chang, S.W., Nguyen, D.D., Zhang, X., Liang, H., Xue, S. (2020). Selective carbon sources and salinities enhance enzymes and extracellular polymeric substances extrusion of *Chlorella* sp. for potential co-metabolism. **Bioresource Technology**, 303, 122877. <https://doi.org/10.1016/j.biortech.2020.122877>
- Wafi, A., Ariadi, H., Muqsith, A., Mahmudi, M., Fadjar, M. (2021). Oxygen consumption of *Litopenaeus vannamei* in intensive ponds based on the dynamic modeling system. **Journal of Aquaculture and Fish Health**, 10(1), 17-24. <https://doi.org/10.20473/jafh.v12i2.32371>.
- Wang, S., Wu, Y., Wang, X. (2016). Heterotrophic cultivation of *Chlorella pyrenoidosa* using sucrose as the sole carbon source by co-culture with *Rhodotorula glutinis*. **Bioresource Technology**, 220, 615-620. <https://doi.org/10.1016/j.biortech.2016.09.010>.
- Wang, L.M., Jiang, B.J., Zhu, W.B., Fu, J.J., Luo, M.K., Liu, W., Dong, Z.J. (2022). The role of melanocortin 1 receptor on melanogenesis pathway in skin color differentiation of red tilapia. **Aquaculture Reports**, 22, 100946. <https://doi.org/10.1016/j.aqrep.2021.100946>
- Wijaya, N.I., Trisyani, N., Sulestiani, A. (2019). Potensi Pengembangan Budidaya Silvofishery Di Area Mangrove Wonorejo Surabaya. **Jurnal Penelitian Hutan dan Konservasi Alam**, 16(2), 173-189. DOI: <https://doi.org/10.20886/jphka.2019.16.2.173-189> (*In Indonesian*)
- Xie, Y., Lu, K., zhao, X., Ma, R., Chen, J., Ho, S.H. (2019). Manipulating Nutritional Conditions and Salinity-Gradient Stress for Enhanced Lutein Production in Marine Microalga *Chlamydomonas* sp. **Biotechnology Journal**, 14(4), 1800380. <https://doi.org/10.1002/biot.201800380>
- Yinding, T., Xiwen, X., Miao, Q., Jingjing, S., Yiyang, Z., wei, Z., Mengzhu, W., Xuejun, W., Yang, Z. (2021). Lake warming intensifies the seasonal pattern of internal nutrient cycling in the eutrophic lake and potential impacts on algal blooms. **Water Research**, 188, 116570. <https://doi.org/10.1016/j.watres.2020.116570>

- Zhang, X., Cvetkovska, M., Morgan-Kiss, R., Huner, N.P.A., Smith, D.R. (2021). Draft genome sequence of the Antarctic green alga *Chlamydomonas* sp. **UWO241**. **iScience**, 24(2), 102084. DOI: 10.1016/j.isci.2021.102084
- Zhang, M., Zhang, Y., Yu, S., Gao, Y., Dong, J., Zhu, W., Wang, X., Li, X., Li, J., Xiong, J. (2023). Two machine learning approaches for predicting cyanobacteria abundance in aquaculture ponds. **Ecotoxicology and Environmental Safety**, 258, 114944. <https://doi.org/10.1016/j.ecoenv.2023.114944>
- Zhou, Y., He, Y., Zhou, Z., Xiao, X., Wang, M., Chen, B. (2022). A newly isolated microalga *Chlamydomonas* sp. YC to efficiently remove ammonium nitrogen of rare earth elements wastewater. **Journal of Environmental Management**, 316, 115284. <https://doi.org/10.1016/j.jenvman.2022.115284>

● **6% Overall Similarity**

Top sources found in the following databases:

- 6% Publications database
- Crossref database
- Crossref Posted Content database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	H Ariadi, M. B Syakirin, S Hidayati, B D. Madusari, H Soeprapto. "Fluctu... Crossref	1%
2	Ihn-Han Bae, Sun-Jin Oh, Stephan Olariu. "Chapter 24 A Fuzzy Logic-Ba... Crossref	<1%
3	Abdul Muqsith, Abdul Wafi, Heri Ariadi. "Peta Tematik Kesesuaian Para... Crossref	<1%
4	N Asiah, S Sukendi, Y Harjoyudanto, J Junianto, A Yustiati. "Water Qual... Crossref	<1%
5	Biao Zhang, Yixuan Bian, Junsheng Chen, Ziyang Zhang, Shengrui Sun, ... Crossref	<1%
6	Wei-Yen Wu, Andy Ying-Guey Fuh. "Rewritable Liquid Crystal Gratings F... Crossref	<1%
7	Heri Ariadi, Muh. Azril, Tholibah Mujtahidah. "Water Quality Fluctuation... Crossref	<1%
8	C.T. Da, T. Lundh, J.E. Lindberg. " Digestibility of dietary components a... Crossref	<1%
9	Atika Marisa Halim, Anna Fauziyah, Lusiana B. Ritonga, Moh. Zainal Ari... Crossref	<1%

- 
- 10 M.M.R. Williams. "A model for the transport of vapour, gas and aerosol... <1%  
Crossref
- 
- 11 Mohammad Mahmudi, Muhammad Musa, Alamanda Bunga, Nur Wati, ... <1%  
Crossref
- 
- 12 A. Guerrero Aguilar, C. Schüth, U. E. Rodriguez Castrejon, B. Noriega Lu... <1%  
Crossref
- 
- 13 Belinda Astari, Irzal Effendi, Tatag Budiardi, Yani Hadiroseyani, Iis Diati... <1%  
Crossref posted content