

PAPER NAME

The abundance of Microcystis sp. on intensive shrimp ponds.pdf

WORD COUNT

3800 Words

CHARACTER COUNT

20539 Characters

PAGE COUNT

6 Pages

FILE SIZE

440.3KB

SUBMISSION DATE

May 15, 2023 7:11 AM GMT+7

REPORT DATE

May 15, 2023 7:12 AM GMT+7

● **9% Overall Similarity**

The combined total of all matches, including overlapping sources, for each database.

- 7% Publications database
- Crossref Posted Content database
- Crossref database
- 8% Submitted Works database

● **Excluded from Similarity Report**

- Internet database
- Quoted material
- Small Matches (Less than 8 words)
- Bibliographic material
- Cited material



The abundance of *Microcystis* sp. on intensive shrimp ponds

Hayati Soeprapto¹, Heri Ariadi^{1,*}, Ubad Badrudin², Petrus Hary Tjahja Soedibya³

² Department of Aquaculture, Faculty of Fisheries, Universitas Pekalongan, Pakalongan, Indonesia.

² Department, of Ariculture, Faculty of Agriculture, Universitas Pekalongan, Pakalongan, Indonesia.

³ Department of Aquaculture, Faculty of Fisheries and Marine Science, Universitas of Jenderal Soedirman, Purwokerto, Indonesia.

ARTICLE INFO

ABSTRACT

Keywords:

Cultivation
Genus
Grazing
Model
Waters

Microcystis sp is plankton species that is detrimental in shrimp pond ecosystems. The purpose of this study was to detect the abundance of plankton and the presence of *Microcystis* sp. in the waters of vaname shrimp (*L. vannamei*) aquaculture ponds. The research method used is descriptive quantitative and qualitative with added analysis of dynamic modeling systems. The results showed that pond water quality conditions were still good enough for shrimp farming activities with details of dissolved oxygen 4.01-6.17 mg/L, pH 7.7-7.9, salinity 20‰, temperature 27.05°-28.25°C, nitrate 0.01-0.15 mg/L, phosphate 0.05-0.20 mg/L, brightness 25-35 cm. For the classification of plankton found among them are from the Bacillariophyceae class, there are the genera *Skeletonema* sp., *Cyclotella* sp., *Coscinodiscus* sp., and *Amphipora* sp., from the Chlorophyceae class, there are the genera *Chlorella* sp., *Oocystis* sp., and *Clamydomonas* sp., from the Cyanophyceae class, the genus *Oscillatoria* sp., *Microcystis* sp., and *Anabaenopsis* sp. The plankton diversity index (H') was obtained between 0.01-0.33 and the uniformity index (E) was obtained between 0.10-0.50, meaning that the condition of the pond waters is still quite good and there are minimal indications of pollution. The abundance of *Microcystis* sp. obtained between 0.2×10^4 - 0.6×10^4 cell/ml. From the results of dynamic modeling analysis it is shown that the abundance of *Microcystis* sp. and waters is influenced by the value of the productivity level of the waters and the process of grazing organisms. The results of this study can be concluded that the plankton abundance rate found in ponds ranges from 0.4×10^4 - 1.9×10^4 cell/ml which consists of 10 genera and 3 classes of plankton. Then *Microcystis* sp. its presence was found in 3 ponds with abundance rate ranging from 0.2×10^4 - 0.6×10^4 cell/ml.

DOI: 10.13170/depik.12.1.30433

Introduction

Shrimp cultivation is one of the many cultivation activities carried out in the Pekalongan coastal. Shrimp cultivation can be carried out in various patterns, such as: traditional, semi-intensive and intensive (Ariadi and Wafi, 2020). The water quality dynamics in shrimp ponds always fluctuate throughout the cultivation cycle (Ariadi et al., 2021). Water quality fluctuations are in line with the intensity level of treatment given to the pond ecosystems (Case et al, 2008).

The dynamics of quality and treatment intensity carried out in ponds will greatly affect the structure and abundance of plankton (Ariadi et al., 2019). Plankton in pond water ecosystems have an

important role as live food and environmental bioindicators (Lyu et al., 2021). The presence of nutrients and temperature are some of the limiting factors that affect the dynamics of plankton dominance in pond waters. The primary productivity rate in pond waters, which is closely related to the plankton abundance, will continue to change, forming character patterns in each cultivation cycle (Zhang et al., 2021).

Plankton several types that are commonly found in vannamei shrimp farming activities in each aquaculture operational cycle are *Microcystis* sp. *Microcystis* sp. is a cyanobacteria that easily blooms and more diverse in pond water ecosystems (Kumar and Sinha, 2014). *Microcystis* sp. has a toxin compound

* Corresponding author.

Email address: ariadi_heri@yahoo.com

called microcystins (Zimba *et al.*, 2006). These microcystins compounds are toxic to shrimp metabolisms. Vaname shrimp is a species of shrimp that is highly resistant to changes in abnormal environmental conditions (Li *et al.*, 2022).

Based on the background above, the purpose of this study was to detect the plankton abundance and the presence of *Microcystis* sp. in the waters of vaname shrimp (*L. vannamei*) aquaculture ponds.

Materials and Methods

Location and time of research

This research was conducted in the shrimp ponds of Siwalan Village on December 2022. The parameters observed in this study included the plankton diversity index, plankton uniformity index, plankton types, and pond water quality parameters. Plankton data collection was carried out in 4 shrimp ponds with the sampling point taking zones being on the surface water, water column and the bottom of the pond.

Data analysis

Data were analyzed descriptively quantitatively and qualitatively using Microsoft Excel and Stella ver.9.0.2 software. Plankton observations were carried out using an Olympus CX23 microscope and Haemocytometer Neubauer©.

Diversity index

The plankton diversity index is calculated based on the Shannon-Wiener index formula by Parsons *et al.* (1977), as follows :

$$H' = \sum_{i=1}^S P_i \log_2 P_i$$

Explanation :

H' : Diversity index

P_i : n_i/N

N_i : Number of individual in- i

N : Total number of individuals

Uniformity index

ndex

The uniformity index is calculated based on the formula introduced by Odum, (1996), as follows :

$$E = H'/H_{\max}$$

Explanation :

E : Species uniformity index

H' : Diversity index

H_{\max} : $\log_2 S$

S : Number of species.

Results

Water quality

The water quality value of shrimp aquaculture ponds at the research location is still quite good. The parameters of dissolved oxygen, salinity, temperature, nitrate, phosphate, and brightness are still in accordance with the quality standard values designated for shrimp farming activities. The water quality values for research ponds can be presented in Table 1. Stable water quality parameter values are possible due to the low level of shrimp stocking density and the minimal chemical treatment given (Amorim and Moura, 2021).

Table 1. Water quality in shrimp pond culture.

Ponds	Water Quality Parameters						
	DO (mg/L)	pH	Salinity (gr/L)	Temperature (°C)	Nitrat (mg/L)	Fosfat (mg/L)	Brightness (cm)
A	4.01-5.67	7.7-7.8	20	27.05-27.75	0.03-0.05	0.05-0.08	30-35
B	5.24-5.83	7.7-7.8	20	27.75-28.00	0.01-0.05	0.07-0.13	30-35
C	4.89-6.17	7.8-7.9	20	27.15-28.25	0.06-0.15	0.13-0.20	25-30
D	5.13-6.04	7.7-7.8	20	27.50-27.75	0.03-0.12	0.06-0.11	30-35

Plankton genus

Based on the results of the species identification analysis, the plankton found in the research pond waters consisted of 10 genera from three classes, namely: Bacillariophyceae (4 genera), Chlorophyceae (3 genera), Cyanophyceae (3 genera). The dominance level of the genus in these waters is low. The plankton genus found in the research pond waters can be seen in (Table 2). the plankton dynamics in research ponds is describe as very diverse as very diverse and

dynamic. Differences in temperature and environmental conditions of pons water ecosystems will trigger plankton dynamics (Munthe *et al.*, 2012).

Based on the available data, it can be shown that the genus bacillariophyceae, chlorophyceae, and cyanophyceae are very dominant compared to other genus. The plankton predominant condition is caused by the ecosystem conditions of the pond is very dynamic (Ariadi *et al.*, 2022). The pond water characters and the intensity of nutrient fluctuations

in pond waters will greatly affect the genus of plankton that grows. Plankton as aquatic microorganisms will easily experience succession if

there are dynamically fluctuations in water conditions (Ariadi *et al.*, 2019). Plankton will easily bloom and mass die if the extreme water.

Table 2. The plankton abundance species found in pond waters.

No.	Class	Plankton abundance in shrimp ponds (cell/ml)			
		A	B	C	D
Bacillariophyceae					
1	<i>Skeletonema</i> sp.	10000			
2	<i>Cyclotella</i> sp.			40000	30000
3	<i>Coscinodiscus</i> sp.				10000
4	<i>Amphipora</i> sp.			10000	
Chlorophyceae					
1	<i>Chlorella</i> sp.	80000	20000		
2	<i>Oocystis</i> sp.	10000		30000	70000
3	<i>Clamydomonas</i> sp.	10000		20000	20000
Cyanophyceae					
1	<i>Oscillatoria</i> sp.	20000	10000		
2	<i>Microcystis</i> sp.	20000		30000	60000
3	<i>Anabaenopsis</i> sp.		10000		
Genera		6	3	5	5
Total (cell/ml)		150000	40000	130000	190000

Diversity index

The plankton diversity index in the study ponds ranged from 0.01-0.33. The highest diversity index value was found in D pond and the lowest in B pond (Figure 1). based on the plankton genus diversity, it can be said that the pond water conditions is still quite good and there are no indications of waste pollution (Munthe *et al.*, 2012). These indications are in accordance with the shrimp culture system conditions at the research location using the traditional plus cultivation system.

The plankton diversity in the shrimp pond will greatly determine the profile of aquatic organisms that grow. The existence of a food chain, especially for higher nekton or shrimp will depend on the presence of primary level organisms, such as plankton and detritus (Amorim *et al.*, 2021). Based on these conditions, the plankton diversity in addition to affecting of water nutrients, also has an impact on the food chain cycle in pond ecosystems. Food chain cycle is natural process can be upgredetable of pond ecosystem (Ariadi, 2020).




Figure 1. Plankton diversity index in pond waters.

Uniformity index

The plankton uniformity index values in ponds ranged between 0.10-0.50, with the highest uniformity index level found in D pond and the lowest in B pond (Figure 2). The average uniformity index value is ≤ 4 , it can be said that the plankton uniformity index level in the study ponds is low (Munthe *et al.*, 2012). The plankton uniformity index will correlate with water quality dynamics and plankton diversity (Amorim and Moura, 2021). In vannamei shrimp ponds, the level of plankton uniformity will continue to change according to the

dynamics of water quality (Ariadi et al., 2019). Water quality in shrimp ponds is a important factor (Wafi et al., 2021).




Figure 2. Plankton uniformity index in ponds.

Microcystis sp. abundance

Total abundance of *Microcystis* sp. in research ponds is dynamic. *Microcystis* sp. almost found in every aquaculture pond except B pond (Figure 3). *Microcystis* sp. is a cosmopolitan plankton that blooms easily under nutrient conditions (Masithah, 2011). Shrimp pond ecosystems that have high nutrient levels are very potential to be used as a living habitat for *Microcystis* sp. (Ariadi et al., 2022).




Figure 1. *Microcystis* sp. abundance in shrimp pond waters.

Based on dynamic modeling analysis, the *Microcystis* sp. will continue to increase of aquatic productivity rate on the waters and the grazing process in the pond ecosystem continues to increase. The growth rate of phototrophic organisms in pond feeding tissue is strongly influenced by the grazing process (Calbet et al., 2012). In shrimp ponds, grazing itself will run naturally under normal conditions (Ariadi et al., 2022). The aquatic productivity rate is proportionally influenced by the fertilization process and the feed waste accumulation (Xiong et al., 2020). The results of dynamic modeling correlation can be presented in (Figure 4).




Figure 4. Results of dynamic modeling system analysis about related prediction *Microcystis* sp. abundance in the shrimp pond.

Discussion

The water quality parameter is an important indicator in the operational cycle of shrimp farming (Madusari et al., 2022). Water quality stabilized will make shrimp comfortable and grow well as well as in the opposite condition (Yu et al., 2022). As an aquatic biota, shrimp are very tolerant of fluctuate change in the conditions of the aquatic ecosystem of their habitat. *L. vannamei* is a shrimp species that is physiologically very strong in resisting the environment fluctuations (Ariadi et al., 2019). The water quality condition is also influenced by the level of treatment carried out during the shrimp culture cycle (Wafi and Ariadi, 2022).

The low level of genus dominance in the research ponds is due to the cultivation system used is the traditional plus model so that not much treatment is given to the cultivation ecosystem. Treatment in the cultivation system will greatly determine the plankton existence (De et al., 2020). The nutrient solubility and water quality stability are also other factors that affect to presence of plankton in ponds (Boyd and Davis, 2020). Erratic weather conditions can also cause the plankton dominance in ponds to be unstable over time (Churnside et al., 2020).

From plankton dominance rate in ponds is related for shrimp stocking density (Aliviyanti et al., 2017). The minimum amount of contaminants can be caused by the low stocking density and shrimp biomass in the ponds. The low stocking density will affect the intensity of nutrients in pond waters (Huang et al., 2020). Plankton will be very sensitive to changes in the aquatic environment dynamics.

The low value of the uniformity index illustrates that the existence of plankton communities in pond water ecosystems tends to be unstable. The dynamic ecosystem of pond waters will affect the level of existence of the plankton (Ariadi and Puspitasari, 2021). Plankton will be very sensitive to changes in environment conditions and food chains that exist in pond aquatic ecosystems (McQuatters-Gollop et al.,

2019). Seeing the low plankton uniformity index value means that the dominance level of plankton is still quite low.

Microcystis sp. is a very tolerant of growing in high-salinity water (Soedarsono et al., 2013). The status of pond waters that have optimum salinity levels and optimal temperature conditions will allow plankton to grow optimally (Ariadi and Mujtahidah, 2022). *Microcystis* sp. in pond waters has a high level of dominance because of its cosmopolitan nature. In trophic waters several types of plankton such as *Oscillatoria* sp., *Microcystis* sp. and *Anabaena* sp. prone to periodic blooming (Aliviayanti et al., 2017).

Based on the results analysis of the diversity index and uniformity index, it was shown that the waters in the research ponds were still quite good. This status correlates closely with the water quality profile and plankton abundance level in pond waters. The dominance and abundance of plankton will follow by water quality dynamics in the waters (McQuatters-Gollop et al., 2019). The plankton dominance and abundance level is also greatly influenced by aquatic productivity rate (Xiong et al., 2020).

Overall, the study results indicate that the condition of pond waters which is still classified as good is closely related with plankton abundance stabilized. The *Microcystis* sp. appearance with primary productivity rate in each research pond. *Microcystis* sp. is a plankton that tends to be adaptive with water condition changes (Huang et al., 2014). Then indirectly, *Microcystis* sp. abundance will greatly affect the grazing process and the food chain in the pond ecosystem

Conclusion

The results of this study can be concluded that the plankton abundance rate found in ponds ranges from 0.4×10^4 - 1.9×10^4 cell/ml which consists of 10 genera and 3 classes of plankton. Then *Microcystis* sp. its presence was found in 3 ponds with abundance rate ranging from 0.2×10^4 - 0.6×10^4 cell/ml.

Acknowledgments

The author would like to thank the Research and Community Service Department of Pekalongan University for grant funding the excellent research programs provided through SK No. 570/B.06.01/LPPM/XI/2022.

References

Aliviayanti, D., Suharjono, C. Retnaningdyah. 2017. Cyanobacteria Community Dynamics and Trophic Status of Intensive Shrimp (*Litopenaeus vannamei*) Farming Pond in Situbondo,

East Java Indonesia. The Journal Of Tropical Life Science, 7(3): 251–257.

- Amorim, C.A., and A.d.N. Moura. 2021. Ecological impacts of freshwater algal blooms on water quality, plankton biodiversity, structure, and ecosystem functioning. Science of The Total Environment, 758: 143605.
- Ariadi, H. 2020. Oksigen Terlarut dan Siklus Ilmiah Pada Tambak Intensif. Gupedia, Bogor.
- Ariadi, H., dan T. Mujtahidah. 2022. Analisis Permodelan Dinamis Kelimpahan Bakteri *Vibrio* sp. Pada Budidaya Udang Vaname, *Litopenaeus vannamei*. Jurnal Riset Akuakultur, 16(4): 255-262.
- Ariadi, H., and M.N. Puspitasari. 2021. Perbandingan Pola Kelayakan Ekologis Dan Finansial Usaha Pada Kegiatan Budidaya Udang Vaname (*L. vannamei*). Fish Scientiae, 11(2): 125-138.
- Ariadi, H., and A. Wafi. 2020. Water Quality Relationship with FCR Value in Intensive Shrimp Culture of Vannamei (*Litopenaeus vannamei*). Samakia: Jurnal Ilmu Perikanan, 11(1): 44-50.
- Ariadi, H., M. Mahmudi, M. Fadjar. 2019. Correlation between density of vibrio bacteria with *Oscillatoria* sp. abundance on intensive *Litopenaeus vannamei* shrimp ponds. Research Journal of Life Science, 6(2): 114-129.
- Ariadi, H., M. Fadjar, M. Mahmudi, Supriatna. 2019. The relationships between water quality parameters and the growth rate of white shrimp (*Litopenaeus vannamei*) in intensive ponds. Aquaculture, Aquarium, Conservation & Legislation, 12(6): 2103-2116.
- Ariadi, H., A. Wafi, B.D. Madusari. 2021. Dinamika Oksigen Terlarut (Studi Kasus Pada Budidaya Udang). Penerbit ADAB, Indramayu.
- Ariadi, H., M.B. Syakirin, S. Hidayati, B.D. Madusari, H. Soeprapto. 2022. Fluctuation Effect of Dissolved of TAN (Total Ammonia Nitrogen) on Diatom Abundance in Intensive Shrimp Culture Ponds. IOP Conference Series: Earth and Environmental Science in Pekanbaru, Indonesia, 2022. IOP Publishing, Pekanbaru Indonesia, pp. 012001.
- Ariadi, H., A. Khristanto, H. Soeprapto, D. Kumalasari, J.L. Sihombing. 2022. Plankton and its potential utilization for climate resilient fish culture. AACL Bioflux, 15(4): 2041-2051.
- Boyd, C.E., dan R.P. Davis. 2020. Lentic Freshwater: Ponds. Aquaculture Ponds, 4: 316-324.
- Calbet, A., R.A. Martinez, S. Isari, S. Zervoudaki, J.C. Nejstgaard, P. Pitta, A.F. Sazhin, D. Sousoni, A. Gomes, S.A. Berger, T.M. Tsagaraki, R. Ptacnik. 2012. Effects of light availability on mixotrophy and microzooplankton grazing in an oligotrophic plankton food web: Evidences from a mesocosm study in Eastern Mediterranean waters. Journal of Experimental Marine Biology and Ecology, 424–425: 66-77.
- Case, M., E.E. Leca, S.N. Leita, E.E. Sant'Anna, R. Schwaborn, A.T. de Moraes Jr. 2008. Plankton community as an indicator of water quality in tropical shrimp culture ponds. Marine Pollution Bulletin, 56: 1343–1352.
- Churnside, J.H., R.D. Marchbanks, S. Vagle, S.W. Bell, P.J. Staben. 2020. Stratification, plankton layers, and mixing measured by airborne lidar in the Chukchi and Beaufort seas. Deep-Sea Research Part II, 177: 104742.
- De, D., K.P. Sandeep, S. Kumar, R.A. Raja, P. Mahalakshmi, T. Sivaramakrishnan, K. Ambasanar, K.K. Vijayan. 2020. Effect of fish waste hydrolysate on growth, survival, health of *Penaeus vannamei* and plankton diversity in culture systems. Aquaculture, 524: 735240.
- Huang, Y., Y. Bai, Y. Wang, H. Kong. 2014. *Solidago canadensis* L. extracts to control algal (*Microcystis*) blooms in ponds. Ecological Engineering, 70: 263-267.
- H Huang, Q., S. Olenin, L. Li, S. Sun, M.D. Troch. 2020. Meiobenthos as food for farmed shrimps in the earthen ponds: Implications for sustainable feeding. Aquaculture, 521: 735094.
- Kumar, B., dan A. Sinha. 2014. Microcystis Toxic Blooms In Fish Culture Ponds And Their Biological And Chemical Control.

- International Journal Of Scientific & Technology Research, 3(3): 398-410.
- Li, X., Y. Chen, X. Chen, S. Zhang, X. Dong, S. Chi, J. Deng, B. Tan, S. Xie. 2022. Cholesterol supplementation improved growth performance, cholesterol metabolism, and intestinal health of Pacific white shrimp (*Litopenaeus vannamei*) fed a low fishmeal diet. *Aquaculture Reports*, 27: 101351.
- Lyu, T., W. Yang, H. Cai, J. Wang, Z. Zheng, J. Zhu. 2021. Phytoplankton community dynamics as a metrics of shrimp healthy farming under intensive cultivation. *Aquaculture Reports*, 21: 100965.
- Madusari, B.D., H. Ariadi, D. Mardhiyana. 2022. Effect of the feeding rate practice on the white shrimp (*Litopenaeus vannamei*) cultivation activities. *Aquaculture, Aquarium, Conservation & Legislation-International Journal of the Bioflux Society*, 15(1): 473-479.
- Masithah, E.D. 2011. Upaya Menurunkan Dominansi *Microcystis aeruginosa* Menggunakan Enzim Pektinase Dari *Pseudomonas pseudomallei*. *Berkala Penelelitian Hayati Edisi Khusus*, 4: 83–86.
- McQuatters-Gollop, A., A. Atkinson, A. Aubert, J. Bedford, M. Best, E. Bresnan, K. Cook, M. Devlin, R. Gowen, D.G. Johns, M. Machairopoulou, A. McKinney, A. Mellor, C. Ostle, C. Scherer, P. Tett. 2019. Plankton lifeforms as a biodiversity indicator for regional-scale assessment of pelagic habitats for policy. *Ecological Indicator*, 101: 913-925.
- Munthe, Y.V., R. Aryawati, Isnaini. 2012. Struktur Komunitas dan Sebaran Fitoplankton di Perairan Sungsang Sumatera Selatan. *Maspari Journal*, 4(1): 122-130.
- Rakhmanda, A., A. Pribadi, Parjiyo, B.I.G. Wibisono. 2021. Production performance of white shrimp *Litopenaeus vannamei* with super-intensive culture on different rearing densities. *Jurnal Akuakultur Indonesia*, 20(1): 56–64.
- Soedarsono, P., S. Rudiyaniti, N. Sukmawati. 2013. Analisis Perbandingan Fitoplankton Dominan Pada Peningkatan Salinitas Dalam Tahapan Pembuatan Garam Dan Kultur Skala Laboratorium. *Journal Of Management Of Aquatic Resources*, 2(3): 1-10.
- Wafi, A., H. Ariadi, A. Muqsith, M. Mahmudi, M. Fadjar. 2021. Oxygen consumption of *Litopenaeus vannamei* in intensive ponds based on the dynamic modeling system. *Journal of Aquaculture and Fish Health*, 10(1): 17-24.
- Wafi, A., and H. Ariadi. 2022. Estimasi Daya Listrik Untuk Produksi Oksigen Oleh Kincir Air Selama Periode “Blind Feeding” Budidaya Udang Vaname (*Litopenaeus vannamei*). *Saintek Perikanan: Indonesian Journal of Fisheries Science and Technology*, 18(1): 19-35.
- Xiong, W., X. Mei, X. Meng, H. Chen, H. Yang. 2020. Phytoplankton biomarkers in surface sediments from Liaodong Bay and their potential as indicators of primary productivity. *Marine Pollution Bulletin*, 159: 111536.
- Yu, P., H. Sahn, Y. Cheng, J. Ma, K. Wang, H. Li. 2022. Translucent disease outbreak in *Penaes vannamei* post-larva accompanies the imbalance of pond water and shrimp gut microbiota homeostasis. *Aquaculture Reports*, 27: 101410.
- Zhang, M., J. Dong, Y. Gao, Y. Liu, C. Zhou, X. Meng, X. Li, M. Li, Y. Wang, D. Dai. 2021. Patterns of phytoplankton community structure and diversity in aquaculture ponds, Henan, China. *Aquaculture*, 544: 737078.
- Zimba, P.V., A. Camus, E.H. Allen, J.M. Burkholder. 2006. Co-occurrence of white shrimp, *Litopenaeus vannamei*, mortalities and microcystin toxin in a southeastern USA shrimp facility. *Aquaculture*, 261: 1048–1055.

How to cite this paper:

Soeprapto, H., H. Ariadi, U. Badrudin, P.H.T. Soedibya. 2023. The abundance of *Microcystis sp.* on intensive shrimp ponds. *Depik Jurnal Ilmu-Ilmu Perairan, Pesisir dan Perikanan*, 12(1): 105-110.

● 9% Overall Similarity

Top sources found in the following databases:

- 7% Publications database
- Crossref database
- Crossref Posted Content database
- 8% Submitted Works database

TOP SOURCES

The sources with the highest number of matches within the submission. Overlapping sources will not be displayed.

1	Universitas Diponegoro on 2020-07-30	2%
	Submitted works	
2	fpptijateng on 2022-02-09	<1%
	Submitted works	
3	Abdul Wafi, Rahwan, Hayati Soeprapto. "POTENSI HASIL TANGKAPAN...	<1%
	Crossref	
4	Heri Ariadi, Tholibah Mujtahidah. "ANALISIS PERMODELAN DINAMIS K...	<1%
	Crossref	
5	S Elviana, S Sunarni, M R Maturbongs, Sajriawati, S Fakhriyyah. "Muds...	<1%
	Crossref	
6	Universitas Khairun on 2022-02-04	<1%
	Submitted works	
7	Syiah Kuala University on 2022-10-29	<1%
	Submitted works	
8	Forum Perpustakaan Perguruan Tinggi Indonesia Jawa Timur II on 202...	<1%
	Submitted works	
9	Pohang University of Science and Technology (POSTECH) on 2022-0...	<1%
	Submitted works	

-
- 10 Vivy Melati, Luthfiana Aprilianita Sari, Yudi Cahyoko, Sulastri Arsad, Ku... <1%
Crossref
-
- 11 Forum Perpustakaan Perguruan Tinggi Indonesia Jawa Timur II on 202... <1%
Submitted works
-
- 12 Tianying Lyu, Wen Yang, Huifeng Cai, Jianping Wang, Zhongming Zhen... <1%
Crossref
-
- 13 Forum Perpustakaan Perguruan Tinggi Indonesia Jawa Timur II on 202... <1%
Submitted works
-
- 14 Johan Risandi, Vivi Yovita Indriasari, Rudhy Akhwady, Agus Sufyan. "R... <1%
Crossref
-
- 15 Syiah Kuala University on 2019-03-23 <1%
Submitted works
-
- 16 H Ariadi, M. B Syakirin, S Hidayati, B D. Madusari, H Soeprapto. "Fluctu... <1%
Crossref